2019

ZOOLOGY

(Honours)

Paper: ZOO-HC-1016

(Theory)

Full Marks: 60

Time: 3 hours

The figures in the margin indicate full marks for the questions

- 1. Choose the correct answer (any seven): 1×7=7
 - (a) Which of the following belongs to Anthozoa?
 - (i) Aurelia
 - (ii) Fungia
 - (iii) Stercularia
 - (iv) Dugesia

- (b) Which of the following does not have any alimentary canal?
 - (i) Taenia
 - (ii) Frog
 - (iii) Earthworm
 - (iv) Ascaris
- (c) The infective stage of Entamoeba histolytica is
 - (i) sporozoite
 - (ii) minuta form
 - (iii) mature cyst
 - (iv) trophic form
- (d) Classification of phylum Porifera mainly based on
 - (i) canal system
 - (ii) spicules
 - (iii) shape of choanocytes
 - (iv) archaeocytes

- (e) Animals devoid of respiratory, excretory and circulatory organs are
 - (i) liver fluke
 - (ii) tapeworms
 - (iii) threadworms
 - (iv) sponges
- (f) In flatworms, the excretory organs are
 - (i) archaeocytes
 - (ii) solenocytes
 - (iii) nephrons
 - (iv) nephridia
- (g) Entamoeba histolytica differs from Amoeba proteus due to absence of
 - (i) contractile vacuole
 - (ii) pseudopodia
 - (iii) binary fission
 - (iv) multiple fission
- (h) Secondary host of Taenia solium is
 - (i) cow
 - (ii) man
 - (iii) sheep
 - (iv) pig

- (i) The body of sponges is mainly composed of
 - (i) spongin fibres
 - (ii) mesoglea
 - (iii) spicules
 - (iv) nematoblasts
- 2. Match the following Column—I with Column—II (any four): 2×4=8
 - (a) Column—I Column—II
 - (i) Coller cell (1) Amphiblastula
 - (ii) Sycon (2) Glass rope sponge
 - (iii) Hyalonema (3) Storage cells
 - (iv) Theocytes (4) Choanocytes
 - (b) Column—I Column—II
 - (i) Cilia (1) Flagella
 - (ii) Minuta form (2) Plasmodium
 - (iii) Signet ring (3) Entamoeba
 - (iv) Euglena (4) Paramoecium
 - (c) Column—I Column—II
 - (i) Ctenophora (1) Limnea
 - (ii) Obelia (2) Gammule
 - (iii) Fasciola (3) Medusa
 - (iv) Freshwater (4) Swimming plates sponges

- (d) Column—I Column—II
 - (i) Euglena (1) Medusa
 - (ii) Ctenophores (2) Offense and defense
 - (iii) Obelia (3) Photosynthetic protist
 - (iv) Dactylozooid (4) Hermaphrodite
- (e) Column—I Column—II
 - (i) Anthozoa (1) Radial or biradial
 - (ii) Hydrozoa (2) Medusoid
 - (iii) Scyphozoa (3) Polypoid
 - (iv) Cnidaria (4) Pennatula
- (f) Column—I Column—II
 - (i) Statocyst (1) Coral formation
 - (ii) Ctenophora (2) Skeleton of a solitary coral
 - (iii) Corallite (3) Sense organ
 - (iv) Millepora (4) Triploblastic origin of tissue
- 3. Answer any three from the following questions: 5×3=15
 - (a) Classify the phylum Nemathelminthes with general characters up to class and give examples.
 - (b) Write about the evolutionary significance of Ctenophora.

- (c) Discuss about the different types of locomotory organs in Protista. Add a note on their significance.
- (d) Write about the pathogenicity of Wuchereria bancrofti.
- (e) Discuss about the different types of spicules of sponges with necessary diagram.
- 4. Answer any three from the following: 10×3=30
 - (a) Discuss the life history of Plasmodium
 vivax with necessary diagrams. Add a
 note on its pathogenicity. 8+2=10
 - (b) Describe the canal system in Porifera with necessary illustrations.
 - (c) Write an essay on the evolution of symmetry and segmentation of Metazoa with necessary illustrations.
 - (d) Discuss the life cycle of Taenia solium with necessary diagrams.

- (e) Write short notes on any two from the following: 5×2=10
 - (i) Parasitic adaptation in helminths
 - (ii) Metagenesis in obelia
 - (iii) Corals and coral reefs
- (f) Write an essay on polymorphism in Cnidaria. 10
